

# Integration by parts

We begin our journey by talking about a trick that could be seen as the reverse for the product rule for derivatives. We will use it to develop an entirely new way to solve problems by minimizing a function of functions. Finally, this will follow Wikipedia contributors (2023), which has the classical derivation, with some pieces added by me.

## Sketch for deriving the integration-by-parts formula

We will use the  $'$  symbol to indicate a derivative with respect to  $x$ . If you know the basics of derivation, you recall that:

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x)$$

Now, let's integrate on both sides:

$$\begin{aligned}\int (f(x)g(x))' dx &= \int f'(x)g(x)dx + \int f(x)g'(x)dx \\ f(x)g(x) &= \int f'(x)g(x)dx + \int f(x)g'(x)dx\end{aligned}$$

If we rename  $u = f(x)$  and  $dv = g'(x)dx$ , then we get the familiar expressions for indefinite and definite integrals:

$$\begin{aligned}\int u dv &= uv - \int v du \\ \int_a^b u dv &= uv|_a^b - \int_a^b v du\end{aligned}$$

Finally, imagine that the functions vanish on the extremes  $a$  and  $b$ . That is,  $u(b)v(b) - u(a)v(a) = 0$ . In that case, you can simplify even further:

$$\int u dv = - \int v du$$

## Calculus of variations

Malliavin Calculus is sometimes called an extension of calculus of variations to stochastic processes. What does that even mean? Let's deal with the first part.

### Derivation and application

I'll assume that you know the basics of calculus. That means you can solve problems like this: let's say we have a function  $f$  and we want to know the value of  $x$  where  $f$  attains its minimum:

$$\min f(x) = x^2 - 4x + 5$$

To solve it, we take the derivative, make it equal to zero, and obtain the arguments that make it so.

$$\begin{aligned} f'(x) &= 2x - 4 = 0 \\ 2x &= 4 \\ x &= 2 \end{aligned}$$

Which means that  $\min f(x) = f(2) = 2^2 - 4 * 2 + 5 = 1$ .

This approach works great when you try to minimize a function **value** over the reals,  $f : \mathbb{R} \rightarrow \mathbb{R}$ , or in general for multivariate functions,  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ .

But let's say we want to find the **function**  $y = f(x)$  that minimizes an expression. We don't know how that function looks, or want to impose any preconceived notions like being a polynomial or a sum of sine/cosine pairs. What do we do, then?

From now on, it's autopilot from any course (I'll be using the Wikipedia article as the baseline). The first step is to define a "function of functions" as the starting point for minimization, and that function is essentially the integral like this:

$$J[y] = \int_{x_1}^{x_2} L(x, y, y') dx$$

To keep things grounded on reality, we will use the classical example of finding the function that represents the smallest distance between two points,  $a$  and  $b$ . Starting with a "coarse grain" case with deltas, we are minimizing the sum of all the little changes in the path between those two points using Pythagoras:

$$\min \sum_{i=1}^n \sqrt{(\Delta x_i)^2 + (\Delta y_i)^2}$$

As little changes become infinitesimal, and with some extreme abuse of notation, we switch to an integral and obtain the expression for  $L$ :

$$\begin{aligned} \int_a^b \sqrt{dx^2 + dy^2} &= \int_a^b \sqrt{(dx)^2 \left( 1 + \left( \frac{dy}{dx} \right)^2 \right)} \\ &= \int_a^b \sqrt{1 + (y')^2} \sqrt{(dx)^2} \\ &= \int_a^b \sqrt{1 + (y')^2} dx \\ &= \int_a^b L(y') dx \end{aligned}$$

Now, we will introduce a generic function  $\eta(x)$  called the variation. This function represents a perturbation and the only requirement that we will impose on it is that  $\eta(a) = \eta(b) = 0$ . We can assume that  $f(x) = y$  is the solution and  $\eta$  is the deviation from the solution, which we can multiply by a small number  $\varepsilon$ . We can then define a function over  $\varepsilon$ ,  $\Phi(\varepsilon) = J[f + \varepsilon * \eta]$ , so that  $\Phi(0) = f$ , our solution.

The trick here is that now, instead of minimizing something that depends on unknown functions, which we can't even start to comprehend, we minimize something that depends on a number,  $\varepsilon$ . Not only that, we constructed  $\varepsilon$  and  $\eta$  so that we know, in advance, that the function we are looking for, the solution, is at  $\Phi(\varepsilon = 0) = J[f]$  and  $\Phi'(0) = 0$  because it's the minimum. Absolute genius.

$$\Phi'(0) = 0 = \left. \frac{d\Phi}{d\varepsilon} \right|_{\varepsilon=0} = \int_a^b \left. \frac{dL}{d\varepsilon} \right|_{\varepsilon=0} dx$$

We take the total derivative of  $\Phi$  over  $\varepsilon$ , just like before, but now  $y = f + \varepsilon\eta$  and  $y' = f' + \varepsilon\eta'$ :

$$\begin{aligned} \frac{dL}{d\varepsilon} &= \frac{\partial L}{\partial x} \underbrace{\frac{dx}{d\varepsilon}}_{=0} + \frac{\partial L}{\partial y} \underbrace{\frac{dy}{d\varepsilon}}_{=\eta} + \frac{\partial L}{\partial y'} \underbrace{\frac{dy'}{d\varepsilon}}_{=\eta'} \\ &= \frac{\partial L}{\partial y} \eta + \frac{\partial L}{\partial y'} \eta' \end{aligned}$$

This looks meaningless, but now comes the magic:

$$\begin{aligned}
\int_a^b \frac{dL}{d\varepsilon} \Big|_{\varepsilon=0} dx &= \int_a^b \left( \frac{\partial L}{\partial f} \eta + \frac{\partial L}{\partial f'} \eta' \right) dx \\
&= \int_a^b \frac{\partial L}{\partial f} \eta dx + \underbrace{\frac{\partial L}{\partial f'} \eta \Big|_a^b}_{=0} - \int_a^b \eta \frac{d}{dx} \frac{\partial L}{\partial f'} dx \\
&= \int_a^b \left( \frac{\partial L}{\partial f} \eta - \eta \frac{d}{dx} \frac{\partial L}{\partial f'} \right) dx \\
0 &= \int_a^b \eta(x) \left( \frac{\partial L}{\partial f} - \frac{d}{dx} \frac{\partial L}{\partial f'} \right) dx
\end{aligned}$$

We start from  $\Phi'(0)$  and we do an *integration by parts*. This is crucial, without that operation this whole scheme falls apart. We also know that  $\eta(a) = \eta(b) = 0$ , so that term vanishes. Joining terms and taking a common factor we reach our final conclusion. The final piece of magic is realizing that this integral is zero, no matter what  $\eta(x)$  we pick. By the Fundamental Lemma of Calculus of Variations, that leaves only one possibility, and it's so important that it's called the Euler-Lagrange equation:

$$\frac{\partial L}{\partial f} - \frac{d}{dx} \frac{\partial L}{\partial f'} = 0$$

Whatever the solution function  $f$  is, it must fulfill that condition.

### Shortest Path Example

Why is it helpful? Let's return to the shortest path between two points, and I'm using  $f = y$  interchangeably. When we left off, we had arrived at:

$$L(f') = \sqrt{1 + (f')^2}$$

We apply the Euler-Lagrange equation and we see where it leads us. What makes this example super clean is that there's no explicit  $f$  in it, only  $f'$ , so we end up with a very small formula:

$$\begin{aligned}
\frac{\partial L}{\partial f} - \frac{d}{dx} \frac{\partial L}{\partial f'} &= 0 \\
-\frac{d}{dx} \frac{f'(x)}{\sqrt{1 + [f'(x)]^2}} &= 0
\end{aligned}$$

If the derivative is zero, then we can integrate that and the result is an unknown constant  $c < 1$ . We do some minor algebra, plus an integration at the end, and we are done!

$$\begin{aligned} \frac{f'(x)}{\sqrt{1 + [f'(x)]^2}} &= c \\ \frac{[f'(x)]^2}{1 + [f'(x)]^2} &= c^2 \\ [f'(x)]^2 &= c^2 + c^2[f'(x)]^2 \\ [f'(x)]^2 &= \frac{c^2}{1 - c^2} \\ f'(x) &= \sqrt{\frac{c^2}{1 - c^2}} = m \\ f(x) &= mx + b \end{aligned}$$

The function with the shortest path between two points is a straight line, and the only thing we needed to know is how the distance between two points is calculated to get a closed-form, analytic solution.

---

Wikipedia contributors. 2023. "Calculus of Variations — Wikipedia, the Free Encyclopedia." [https://en.wikipedia.org/wiki/Calculus\\_of\\_variations](https://en.wikipedia.org/wiki/Calculus_of_variations).