

Malliavin Derivative

Now that we have established that Cameron-Martin directions allow us to change the integrals under Gaussian measures, we are ready to define the Malliavin derivative. We can do it from two angles:

- We can talk about polynomial chaos, and then arrive to the derivative
- We define the derivative, and not refer to polynomial chaos at all

We will follow the second approach for pedagogical reasons, collecting all the things we have learned. After that, we will follow Friz (2002) and Alos (2021) to bring all down to Earth.

Malliavin Derivative building blocks

We have accumulated a ton of knowledge. It's time for us to approach the point of this entire webpage. When we asked the Question, we mentioned that the derivative of a random variable that follows a Brownian motion with respect to time can't be calculated, that was the whole point: what kind of derivative can we calculate?

We will consider a Cameron-Martin direction of $\hat{h}(\tau) = \int_0^\tau h dt$. Moreover, $\hat{h} \in \hat{H}$, the Cameron-Martin space. This definition allow us to have a derivative over time for the direction, so $\frac{\partial \hat{h}(\tau)}{\partial \tau} = h$, with $h \in H = L^2[0, 1]$, the space of functions that are continuous between 0 and 1, and that allow a 2-norm¹.

Brownian motion paths up to τ will be denoted as $\omega(\tau)$. The function $1_{[0, \tau]}$ is a function that's 1 in the indicated interval and 0 everywhere else, behaving like a filter. Finally, we will

¹On a more formal write-up, we can consider this translation:

$$T_{\hat{h}}(\omega) = \omega + \varepsilon \int_0^\tau h = \omega + \varepsilon \hat{h}$$

And the change of measure/variable, that is, our Radon-Nikodym derivative, will be:

$$\frac{d(T_{\hat{h}})_* W}{dW}(\omega) = \exp \left(\int_0^1 h d\beta(\omega) - \frac{1}{2} \int_0^1 h^2 d\lambda \right)$$

Finally, without loss of generality, we can assume $\|h\| = 1$, and let all the h_i be orthonormal in H . This is a safe assumption due to Gram-Schmidt orthonormalization.

define a function like $W(1_{[0,\tau]})$ as an Ito integral of the filter . We will also use B_t and dB_t to reference Brownian motions, alongside W_t and dW_t .

We will also consider a polynomial function² f that takes n Brownian motions as variables, each stopping at a different time. Two Brownian motions from 0 to t are essentially the same, even if their paths are different. So, each Brownian motion will go from 0 to t_1, t_2, \dots, t_n . This function f is for the moment pretty useless and basic. We will find its use and extensions later.

Now, we will do something just like in the section for calculus of variations. Let's have $f(\omega + \varepsilon \hat{h})$. This means that we are moving each of the n Brownian motion path a bit along a Cameron-Martin direction, so all is good. Finally, we will take the derivative with respect to ε , and we will evaluate the derivative at $\varepsilon = 0$. The Brownian motions, which can't be derived, won't be touched because they don't depend on ε .

And here's the final result:

$$\begin{aligned} \left. \frac{d}{d\varepsilon} F(\omega + \varepsilon \hat{h}) \right|_{\varepsilon=0} &= \left[\sum_{i=1}^n \partial_i f(\omega(t_1) + \varepsilon \hat{h}, \omega(t_2) + \varepsilon \hat{h}, \dots, \omega(t_n) + \varepsilon \hat{h}) \int_0^{t_i} h d\lambda \right]_{\varepsilon=0} \\ &= \sum_{i=1}^n \partial_i f(\omega(t_1), \omega(t_2), \dots, \omega(t_n)) \int_0^{t_i} h d\lambda \\ &= \langle DF, h \rangle_H \end{aligned}$$

We define DF as:

$$DF = \sum_i \partial_i f(W(1_{[0,\tau_1]}), \dots, W(1_{[0,\tau_n]})) 1_{[0,\tau_i]}$$

Or its alternative extension:

$$DF = \sum_i \partial_i f(W(h_1), \dots, W(h_n)) h_i$$

making DF an H -valued random variable.

²A function with polynomial growth is also acceptable. Conditions for this function belong to the theory and don't bring pedagogical clarity.

What did just happen?

This D is the Malliavin Derivative. It is a linear operator (like the ones we saw before) and it's applied to F . It means that if you have a function F with variables composed of these Brownian motions multiplied by these h_i functions, then you can apply this derivative definition.

This F looks very limited right now, but if you could *represent, approximate* or find an *equivalence* between the random variable you are interested in and a function like F , then you could apply the derivative to the equivalence. The theory is concerned with finding those connections, we won't bother to do that here.

At this point, the collection of h_i functions are a theoretical requirement and seem pulled out of thin air. For one, they aren't more arbitrary than the displacement h from the classical derivative definition $\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$. As we move forward, we will select the $h_i \in H$ that will be suitable for our purposes. More theoretical sources may establish that h_i are orthonormal (or be made orthonormal), especially under an inner product with a Gaussian weight. Instead, we will follow the simplest, default choice: we will use the single function $h = 1_{[0,1]}$, that is, a function that's 1 on the interval and 0 elsewhere. This helps when the function is inside an integral because it "cuts" or "filters" the integration domain, like in this example:

$$\int_{-\infty}^{+\infty} f(x, t) \cdot 1_{[0,1]}(x) dx = \int_0^1 f(x, t) dx$$

A second question is whether the $[0, 1]$ interval is a fundamental limit or if we can extend this to at least $[0, T]$, which is what we see in common stochastic processes. All authors focusing on the theoretical aspects of the derivative stay in that interval, seemingly treating 1 as a conventional limit. On the other hand, Alos (2021) extends it. So, from now on, we use $h = 1_{[0,T]}$ or $h = 1_{[0,1]}$ based purely on convenience.

Examples

Now, let's start with a *very* simple example. Let's calculate the Malliavin derivative of a Wiener process/Brownian motion:

$$F = W(h) = \int_0^1 h dB$$

Let h be as we defined above, a function of L^2 in the interval $[0, 1]$. In this case, the Malliavin derivative is:


$$DF = D(W(h)) = h$$

Here's a good rule-of-thumb: it's as if we did $\frac{\partial F}{\partial W} \cdot h$ instead of $\frac{\partial F}{\partial t}$ (which we know that we can't do). We are seeing how much the function changes as an underlying Brownian motion changes (on a Cameron-Martin direction). Here's a plot of how different choices of $h(t)$ and a sample path for $W(h(t))$ look like:


```
library(ggplot2)
library(rgl)

# Setup
steps = 100000
h <- 1/steps
t <- seq(from = 0, to = 1, length.out=steps)
dBt <- c(0, rnorm(n=(steps-1),mean = 0,sd = sqrt(h)))

# Left
h_1 <- Vectorize(function(x) 1)
h_hat_1 <- cumsum(h_1(t) * dBt)
ggplot(mapping = aes(x = t)) +
  xlab('Time') +
  ylab('f(t)') +
  geom_line(mapping = aes(y=h_hat_1, colour="W(h)"),linetype = 2) +
  geom_line(mapping = aes(y = h_1(t), colour = "h = 1"), linewidth = 1.25) +
  scale_colour_manual("Functions",values=c("black","red"))
```




```
# Middle
h_2 <- Vectorize(function(x) x)
h_hat_2 <- cumsum(h_2(t) * dBt)
ggplot(mapping = aes(x = t)) +
  xlab('Time') +
  ylab('f(t)') +
  geom_line(mapping = aes(y=h_hat_2, colour="W(h)"),linetype = 2) +
  geom_line(mapping = aes(y = h_2(t), colour = "h = t"), linewidth = 1.25) +
  scale_colour_manual("Functions",values=c("black","red"))
```



```

# Right
h_3 <- Vectorize(function(x) x**2 - 1)
h_hat_3 <- cumsum(h_3(t) * dBt)
ggplot(mapping = aes(x = t)) +
  xlab('Time') +
  ylab('f(t)') +
  geom_line(mapping = aes(y=h_hat_3, colour="W(h)"),linetype = 2) +
  geom_line(mapping = aes(y = h_3(t), colour = "h = t2-1"), linewidth = 1.25) +
  scale_colour_manual("Functions",values=c("black","red"))

```



The simple case above also means that we can rewrite our original definition as

$$DF = \sum_i \partial_i f(W(h_1), \dots, W(h_n)) D(W(h_i))$$

which looks like a normal chain rule. I'm not going to prove it, but the product rule is also exactly like what we expect:

$$D(FG) = FD(G) + GD(F)$$

Now, we will do another simple example, a Brownian motion squared:

$$F(t) = (W_t)^2$$


In this case, we use $h(s) = 1_{[0,t]}(s)$. See that there are actually two time-related variables in place: one is the original t that $F(t)$ refers to, and the second one is the one introduced by the Malliavin derivative, which we will call s . With that in mind, we are ready to provide an answer:

$$D_s F = 2W_t (DW_t) = 2W_t h = 2W_t 1_{[0,t]}(s)$$

There are two things to note here. First of all, the derivative takes non-zero values only when $0 < s < t$, and this makes sense: we don't expect the function F to change beyond the time t where it's evaluated. Secondly, this also means that when we plotted $W(h(t))$ and $h(t)$ together, it was a bit fake because we were assuming $s = t$ for the sake of drawing it. Even so, I think that assumption allows us to see there's a connection between the Malliavin derivative and the "instantaneous variance" or volatility of the process: the closer the derivative is to 0, the perturbations or movements are smaller. I'll plot now a sample path for F and DF , with $t \in [0, 1]$ and $s = t$ (which, again, isn't the full picture, but I find it useful to connect the dots):

```
# Setup
steps = 100000
h <- 1/steps
t <- seq(from = 0, to = 1, length.out=steps)
dWt <- c(0, rnorm(n=(steps-1),mean = 0,sd = sqrt(h)))
Wt <- cumsum(dWt)
f <- Wt**2
Df <- 2*Wt

# Plot
ggplot(mapping = aes(x = t)) +
  xlab('Time') +
  ylab('f(t)') +
  geom_line(mapping = aes(y = f, colour="W(t)2"),linetype = 2) +
  geom_line(mapping = aes(y = Df, colour="DW(s=t)"), linewidth = 0.25) +
  scale_colour_manual("Functions",values=c("black","red"))
```

Skorohod integral & more integration by parts

We mentioned that linear operators come with a companion, adjoint operator. Malliavin derivatives aren't the exception, and given that we defined D with an inner product and an h function to support it as in $\langle DF, h \rangle_H$, we will look for an adjoint operator, which we call δ , so that:

$$\langle DF, h \rangle_H = \langle F, \delta(h) \rangle_{R^n}$$

We can find it by doing a bit of manipulation, taking some shortcuts here and relinquishing some mathematical rigor. From the definition, above, and assuming $h = h_1$, we will take the expected value of the inner product, do integration by parts and some replacements, and we get:

$$\begin{aligned}
\mathbb{E} [\langle DF, h \rangle_H] &= \mathbb{E} \left[\sum_i \partial_i f \langle h_i, h \rangle_H \right] \\
&= \int_{\mathbb{R}^n} \underbrace{\partial_1 f(x)}_{dv} (2\pi)^{-n/2} \underbrace{e^{-\|x\|^2/2}}_u dx \\
&= - \int_{\mathbb{R}^n} f(x) (-x_1) \underbrace{(2\pi)^{-n/2} e^{-\|x\|^2/2} dx}_{\text{Gaussian measure}} \\
&= \int_{\mathbb{R}^n} x_1 f(x) d\mu_1 \\
&= \mathbb{E} [F W(h_1)] \\
&= \mathbb{E} [F W(h)] \\
&= \mathbb{E} \left[F \int_0^1 h dB \right]
\end{aligned}$$

What this means is that the expected value of the random variable DF measured using the function h between 0 and 1 as a ruler, is the same as the expected value of F multiplied by that integral of h on the right. We are implying that the δ operator is that integral. This is called the **Skorohod integral**, and it coincides with an Ito integral for non-anticipative processes³:

$$\delta(h) = \int_0^1 h dB_t$$

By now, you probably have noticed how common it is to flip integrals and derivatives using integration by parts. It's almost like second nature with Malliavin Calculus.

What now?

So, we have finally arrived to a definition of the Malliavin Derivative. It has taken us a while, but now we are in a comfortable position to use it for something!

³Formally, this is either predictable or adapted processes. A predictable process has a value at t known in advance (for example, a non-random function). An adapted process has a value at t known at t (for example, the price of a stock). Non-adapted processes, or anticipative processes, are only known after some time has passed, with their information depending on the future (for example, a hale occurring in 7 days).

Alos, & Lorite, E. 2021. *Malliavin Calculus in Finance: Theory and Practice (1st Ed.)*. 1st ed. Financial Mathematics Series. Chapman; Hall/CRC. <https://doi.org/10.1201/9781003018681>.

Friz, Peter K. 2002. "An Introduction to Malliavin Calculus." In. <https://api.semanticscholar.org/CorpusID:2479628>.